

## The Change of Coordinates Matrix

### Previously

- We found the coordinates of  $p(x) = 6 - 2x + 2x^2$  with respect to two different bases

## The Change of Coordinates Matrix

### Example

Let  $B = \{1 + x - x^2, x + x^2, -x + 3x^2\}$  and  $C = \{1 + x + x^2, 1 - x - 2x^2, 4x\}$ , and let  $p(x) \in P_2$  be such that  $[p(x)]_C = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}$ . Find  $[p(x)]_B$ .

The first step to finding  $[p(x)]_B$  is to find what  $p(x)$  is. Using the given  $C$ -coordinates of  $p(x)$ , this is a straightforward calculation:

$$\begin{aligned} p(x) &= 3(1 + x + x^2) + 2(1 - x - 2x^2) + (4x) \\ &= 5 + 5x - x^2 \end{aligned}$$

Now we simply need to find the  $B$ -coordinates of  $5 + 5x - x^2$ .

That is, we need to find scalars  $t_1$ ,  $t_2$ , and  $t_3$  such that

$$5 + 5x - x^2 = t_1(1 + x - x^2) + t_2(x + x^2) + t_3(-x + 3x^2) = (t_1) + (t_1 + t_2 - t_3)x + (-t_1 + t_2 + 3t_3)x^2$$

Setting the coefficients equal to each other, we see that we are looking for the solution to the following system:

$$\begin{array}{rclcl} t_1 & & & & = 5 \\ t_1 & +t_2 & -t_3 & = 5 \\ -t_1 & +t_2 & +3t_3 & = -1 \end{array}$$

### The Change of Coordinates Matrix

#### Example

Let  $B = \{1 + x - x^2, x + x^2, -x + 3x^2\}$  and  $C = \{1 + x + x^2, 1 - x - 2x^2, 4x\}$ , and let  $p(x) \in P_2$  be such that

$$[p(x)]_C = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}. \text{ Find } [p(x)]_B.$$

To find the solution, we will row reduce its augmented matrix:

$$\begin{aligned} & \left[ \begin{array}{ccc|c} 1 & 0 & 0 & 5 \\ 1 & 1 & -1 & 5 \\ -1 & 1 & 3 & -1 \end{array} \right] \begin{array}{l} R_2 - R_1 \\ R_3 + R_1 \end{array} \sim \left[ \begin{array}{ccc|c} 1 & 0 & 0 & 5 \\ 0 & 1 & -1 & 0 \\ 0 & 1 & 3 & 4 \end{array} \right] \begin{array}{l} \\ R_3 - R_2 \end{array} \\ & \sim \left[ \begin{array}{ccc|c} 1 & 0 & 0 & 5 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 4 & 4 \end{array} \right] \begin{array}{l} \\ \frac{1}{4} R_3 \end{array} \sim \left[ \begin{array}{ccc|c} 1 & 0 & 0 & 5 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 1 & 1 \end{array} \right] \begin{array}{l} \\ R_2 + R_3 \end{array} \\ & \sim \left[ \begin{array}{ccc|c} 1 & 0 & 0 & 5 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \end{array} \right] \end{aligned}$$

And so we see that  $t_1 = 5$ ,  $t_2 = 1$ , and  $t_3 = 1$ . And this means that  $[p(x)]_B = \begin{bmatrix} 5 \\ 1 \\ 1 \end{bmatrix}$ .

### The Change of Coordinates Matrix

#### Theorem 4.4.1

Let  $B$  be a basis for a finite dimensional vector space  $V$ . Then, for any  $\mathbf{x}, \mathbf{y} \in V$  and  $t \in \mathbb{R}$ , we have

$$[t\mathbf{x} + \mathbf{y}]_B = t[\mathbf{x}]_B + [\mathbf{y}]_B$$

#### Proof

Let  $B = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ , let  $[\mathbf{x}]_B = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$ , and let  $[\mathbf{y}]_B = \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix}$ .

Then  $\mathbf{x} = x_1\mathbf{v}_1 + \dots + x_n\mathbf{v}_n$  and  $\mathbf{y} = y_1\mathbf{v}_1 + \dots + y_n\mathbf{v}_n$

Then 
$$\begin{aligned} t\mathbf{x} + \mathbf{y} &= t(x_1\mathbf{v}_1 + \dots + x_n\mathbf{v}_n) + (y_1\mathbf{v}_1 + \dots + y_n\mathbf{v}_n) \\ &= (tx_1\mathbf{v}_1 + \dots + tx_n\mathbf{v}_n) + (y_1\mathbf{v}_1 + \dots + y_n\mathbf{v}_n) \\ &= (tx_1 + y_1)\mathbf{v}_1 + \dots + (tx_n + y_n)\mathbf{v}_n \end{aligned}$$

So we see that the  $B$ -coordinates for  $t\mathbf{x} + \mathbf{y}$  are  $\begin{bmatrix} tx_1 + y_1 \\ \vdots \\ tx_n + y_n \end{bmatrix}$ .

Which means that

$$[t\mathbf{x} + \mathbf{y}]_B = \begin{bmatrix} tx_1 + y_1 \\ \vdots \\ tx_n + y_n \end{bmatrix} = t \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} + \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix} = t[\mathbf{x}]_B + [\mathbf{y}]_B$$

□

### The Change of Coordinates Matrix

So how does this Theorem help us?

Well, suppose we have two bases  $B = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$  and  $C = \{\mathbf{w}_1, \dots, \mathbf{w}_n\}$  for a vector space  $\mathbb{V}$ , and let  $\mathbf{x} \in \mathbb{V}$  be

such that  $[\mathbf{x}]_C = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$ .

To find  $[\mathbf{x}]_B$  we use Theorem 4.4.1 to note the following:

$$\begin{aligned} [\mathbf{x}]_B &= [x_1 \mathbf{w}_1 + \dots + x_n \mathbf{w}_n]_B \\ &= x_1 [\mathbf{w}_1]_B + \dots + x_n [\mathbf{w}_n]_B \\ &= \begin{bmatrix} [\mathbf{w}_1]_B & \dots & [\mathbf{w}_n]_B \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} \\ &= \begin{bmatrix} [\mathbf{w}_1]_B & \dots & [\mathbf{w}_n]_B \end{bmatrix} [\mathbf{x}]_C \end{aligned}$$

This means that to find the  $B$ -coordinates for  $\mathbf{x}$ , we can multiply the  $C$ -coordinates by a matrix whose columns are the  $B$ -coordinates of the vectors in  $C$ .

### The Change of Coordinates Matrix

**Definition:** Let  $B$  and  $C = \{\mathbf{w}_1, \dots, \mathbf{w}_n\}$  both be bases for a vector space  $\mathbb{V}$ . The matrix  $P = \begin{bmatrix} [\mathbf{w}_1]_B & \dots & [\mathbf{w}_n]_B \end{bmatrix}$  is called the **change of coordinates matrix** from  $C$ -coordinates to  $B$ -coordinates, and satisfies

$$[\mathbf{x}]_B = P[\mathbf{x}]_C$$

### The Change of Coordinates Matrix

#### Example

Let  $B = \{1 + x - x^2, x + x^2, -x + 3x^2\}$  and  $C = \{1 + x + x^2, 1 - x - 2x^2, 4x\}$ , and let  $p(x) \in P_2$  be such that

$[p(x)]_C = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}$ . Find the change of coordinates matrix from  $C$ -coordinates to  $B$ -coordinates.

To do this, we need to find  $[1 + x + x^2]_B$ ,  $[1 - x - 2x^2]_B$ , and  $[4x]_B$ .

To find the first  $B$ -coordinates, we need to find scalars  $a_1$ ,  $a_2$ , and  $a_3$  such that

$$1 + x + x^2 = a_1(1 + x - x^2) + a_2(x + x^2) + a_3(-x + 3x^2) = (a_1) + (a_1 + a_2 - a_3)x + (-a_1 + a_2 + 3a_3)x^2$$

which is equivalent to the system

$$\begin{array}{rcl} a_1 & & = 1 \\ a_1 + a_2 - a_3 & = & 1 \\ -a_1 + a_2 + 3a_3 & = & 1 \end{array}$$

For the second  $C$  polynomial, we need to find scalars  $b_1$ ,  $b_2$ , and  $b_3$  such that

$$1 - x - 2x^2 = b_1(1 + x - x^2) + b_2(x + x^2) + b_3(-x + 3x^2) = (b_1) + (b_1 + b_2 - b_3)x + (-b_1 + b_2 + 3b_3)x^2$$

which is equivalent to the system

$$\begin{array}{rcl} b_1 & & = 1 \\ b_1 + b_2 - b_3 & = & -1 \\ -b_1 + b_2 + 3b_3 & = & -2 \end{array}$$

### The Change of Coordinates Matrix

#### Example

Let  $B = \{1 + x - x^2, x + x^2, -x + 3x^2\}$  and  $C = \{1 + x + x^2, 1 - x - 2x^2, 4x\}$ , and let  $p(x) \in P_2$  be such that

$[p(x)]_C = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}$ . Find the change of coordinates matrix from  $C$ -coordinates to  $B$ -coordinates.

For the third polynomial in  $C$ , we need to find scalars  $c_1$ ,  $c_2$ , and  $c_3$  such that

$$4x = c_1(1 + x - x^2) + c_2(x + x^2) + c_3(-x + 3x^2) = (c_1) + (c_1 + c_2 - c_3)x + (-c_1 + c_2 + 3c_3)x^2$$

which is equivalent to the system

$$\begin{array}{rcl} c_1 & & = 0 \\ c_1 + c_2 - c_3 & = & 4 \\ -c_1 + c_2 + 3c_3 & = & 0 \end{array}$$

### The Change of Coordinates Matrix

#### Example

Let  $B = \{1 + x - x^2, x + x^2, -x + 3x^2\}$  and  $C = \{1 + x + x^2, 1 - x - 2x^2, 4x\}$ , and let  $p(x) \in P_2$  be such that

$[p(x)]_C = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}$ . Find the change of coordinates matrix from  $C$ -coordinates to  $B$ -coordinates.

Now, all three of these systems have the same coefficient matrix, so we can solve them simultaneously by row reducing the following triply augmented matrix:

$$\left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 1 & 0 \\ 1 & 1 & -1 & 1 & -1 & 4 \\ -1 & 1 & 3 & 1 & -2 & 0 \end{array} \right]$$

$$\begin{array}{l} R_2 - R_1 \\ R_3 + R_1 \end{array} \sim \left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 1 & 0 \\ 0 & 1 & -1 & 0 & -2 & 4 \\ 0 & 1 & 3 & 2 & -1 & 0 \end{array} \right] \begin{array}{l} R_3 - R_2 \\ \frac{1}{4}R_3 \end{array} \sim \left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 1 & 0 \\ 0 & 1 & -1 & 0 & -2 & 4 \\ 0 & 0 & 4 & 2 & 1 & -4 \end{array} \right] \frac{1}{4}R_3 \sim \left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 1 & 0 \\ 0 & 1 & -1 & 0 & -2 & 4 \\ 0 & 0 & 1 & 1/2 & 1/4 & -1 \end{array} \right]$$

$$R_2 + R_3 \sim \left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 1/2 & -7/4 & 3 \\ 0 & 0 & 1 & 1/2 & 1/4 & -1 \end{array} \right]$$

### The Change of Coordinates Matrix

#### Example

Let  $B = \{1 + x - x^2, x + x^2, -x + 3x^2\}$  and  $C = \{1 + x + x^2, 1 - x - 2x^2, 4x\}$ , and let  $p(x) \in P_2$  be such that

$[p(x)]_C = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}$ . Find the change of coordinates matrix from  $C$ -coordinates to  $B$ -coordinates.

Reading off the first augmented column, we see that  $a_1 = 1$ ,  $a_2 = \frac{1}{2}$ , and  $a_3 = \frac{1}{2}$ , so

$$[1 + x + x^2]_B = \begin{bmatrix} 1 \\ 1/2 \\ 1/2 \end{bmatrix}$$

Reading off the second augmented column, we see that  $b_1 = 1$ ,  $b_2 = -\frac{7}{4}$ , and  $b_3 = \frac{1}{4}$ , so

$$[1 - x - 2x^2]_B = \begin{bmatrix} 1 \\ -7/4 \\ 1/4 \end{bmatrix}$$

And reading off the third augmented column, we see that  $c_1 = 0$ ,  $c_2 = 3$ , and  $c_3 = -1$ , so

$$[4x]_B = \begin{bmatrix} 0 \\ 3 \\ -1 \end{bmatrix}$$

### The Change of Coordinates Matrix

#### Example

Let  $B = \{1 + x - x^2, x + x^2, -x + 3x^2\}$  and  $C = \{1 + x + x^2, 1 - x - 2x^2, 4x\}$ , and let  $p(x) \in P_2$  be such that

$[p(x)]_C = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}$ . Find the change of coordinates matrix from  $C$ -coordinates to  $B$ -coordinates.

And this means that our change of coordinates matrix  $P$  is

$$\begin{bmatrix} 1 & 1 & 0 \\ 1/2 & -7/4 & 3 \\ 1/2 & 1/4 & -1 \end{bmatrix}$$

Notice that  $P$  is the same as the right side of our RREF matrix  $\left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 1/2 & -7/4 & 3 \\ 0 & 0 & 1 & 1/2 & 1/4 & -1 \end{array} \right]$

Note also that

$$P \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 0 \\ 1/2 & -7/4 & 3 \\ 1/2 & 1/4 & -1 \end{bmatrix} \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 + 2 + 0 \\ 3/2 - 7/2 + 3 \\ 3/2 + 1/2 - 1 \end{bmatrix} = \begin{bmatrix} 5 \\ 1 \\ 1 \end{bmatrix}$$

which is the same result we got in the original example.

### The Change of Coordinates Matrix

#### Theorem 4.4.2

Let  $B$  and  $C$  both be bases for a finite-dimensional vector space  $V$ . Let  $P$  be the change of coordinates matrix from  $C$ -coordinates to  $B$ -coordinates. Then  $P$  is invertible and  $P^{-1}$  is the change of coordinates matrix from  $B$ -coordinates to  $C$ -coordinates.

#### Proof

To see that  $P^{-1}$  is the change of coordinates matrix from  $B$ -coordinates to  $C$ -coordinates, note that

$$P^{-1}[x]_B = P^{-1}(P[x]_C) = (P^{-1}P)[x]_C = I[x]_C = [x]_C$$

□

### The Change of Coordinates Matrix

#### Example

Let  $S = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}$  be the standard basis for  $M(2, 2)$ , and let

$$B = \left\{ \begin{bmatrix} 1 & 2 \\ 3 & 1 \end{bmatrix}, \begin{bmatrix} -1 & 0 \\ -1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 8 & -3 \end{bmatrix}, \begin{bmatrix} -1 & 4 \\ 1 & 7 \end{bmatrix} \right\}.$$

Find the change of coordinates matrix  $Q$  from  $B$ -coordinates to  $S$ -coordinates, and find the change of coordinates matrix  $P$  from  $S$ -coordinates to  $B$ -coordinates.

### The Change of Coordinates Matrix

#### Solution

The change of coordinates matrix  $Q$  from  $B$ -coordinates to  $S$ -coordinates is

$$\begin{bmatrix} \begin{bmatrix} 1 & 2 \\ 3 & 1 \end{bmatrix}_S & \begin{bmatrix} -1 & 0 \\ -1 & 2 \end{bmatrix}_S & \begin{bmatrix} 3 & 2 \\ 8 & -3 \end{bmatrix}_S & \begin{bmatrix} -1 & 4 \\ 1 & 7 \end{bmatrix}_S \end{bmatrix}$$

But we can find these coordinates without any calculations:

$$\begin{aligned} \begin{bmatrix} 1 & 2 \\ 3 & 1 \end{bmatrix} &= 1 \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + 2 \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} + 3 \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} + 1 \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \\ \begin{bmatrix} -1 & 0 \\ -1 & 2 \end{bmatrix} &= -1 \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + 0 \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} - 1 \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} + 2 \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \\ \begin{bmatrix} 3 & 2 \\ 8 & -3 \end{bmatrix} &= 3 \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + 2 \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} + 8 \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} - 3 \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \\ \begin{bmatrix} -1 & 4 \\ 1 & 7 \end{bmatrix} &= -1 \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + 4 \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} + 1 \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} + 7 \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \end{aligned}$$

so

$$\begin{bmatrix} 1 & 2 \\ 3 & 1 \end{bmatrix}_S = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 & 0 \\ -1 & 2 \end{bmatrix}_S = \begin{bmatrix} -1 \\ 0 \\ -1 \\ 2 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 8 & -3 \end{bmatrix}_S = \begin{bmatrix} 3 \\ 2 \\ 8 \\ -3 \end{bmatrix}, \begin{bmatrix} -1 & 4 \\ 1 & 7 \end{bmatrix}_S = \begin{bmatrix} -1 \\ 4 \\ 1 \\ 7 \end{bmatrix}$$

and so we see that

$$Q = \begin{bmatrix} 1 & -1 & 3 & -1 \\ 2 & 0 & 2 & 4 \\ 3 & -1 & 8 & 1 \\ 1 & 2 & -3 & 7 \end{bmatrix}$$

### The Change of Coordinates Matrix

#### Solution

To find the change of coordinates matrix  $P$  from  $S$ -coordinates to  $B$ -coordinates, we use Theorem 4.4.2, which tells us that  $P = Q^{-1}$ , and then we use the matrix inverse algorithm to find  $Q^{-1}$ :

$$\begin{aligned}
 & \left[ \begin{array}{cccc|cccc} 1 & -1 & 3 & -1 & 1 & 0 & 0 & 0 \\ 2 & 0 & 2 & 4 & 0 & 1 & 0 & 0 \\ 3 & -1 & 8 & 1 & 0 & 0 & 1 & 0 \\ 1 & 2 & -3 & 7 & 0 & 0 & 0 & 1 \end{array} \right] \begin{array}{l} R_2 - 2R_1 \\ R_3 - 3R_1 \\ R_4 - R_1 \end{array} \sim \left[ \begin{array}{cccc|cccc} 1 & -1 & 3 & -1 & 1 & 0 & 0 & 0 \\ 0 & 2 & -4 & 6 & -2 & 1 & 0 & 0 \\ 0 & 2 & -1 & 4 & -3 & 0 & 1 & 0 \\ 0 & 3 & -6 & 8 & -1 & 0 & 0 & 1 \end{array} \right] \begin{array}{l} \\ \frac{1}{2}R_2 \\ \\ \end{array} \\
 & \sim \left[ \begin{array}{cccc|cccc} 1 & -1 & 3 & -1 & 1 & 0 & 0 & 0 \\ 0 & 1 & -2 & 3 & -1 & 1/2 & 0 & 0 \\ 0 & 2 & -1 & 4 & -3 & 0 & 1 & 0 \\ 0 & 3 & -6 & 8 & -1 & 0 & 0 & 1 \end{array} \right] \begin{array}{l} \\ R_3 - 2R_2 \\ R_4 - 3R_2 \end{array} \sim \left[ \begin{array}{cccc|cccc} 1 & -1 & 3 & -1 & 1 & 0 & 0 & 0 \\ 0 & 1 & -2 & 3 & -1 & 1/2 & 0 & 0 \\ 0 & 0 & 3 & -2 & -1 & -1 & 1 & 0 \\ 0 & 0 & 0 & -1 & 2 & -3/2 & 0 & 1 \end{array} \right] \begin{array}{l} \\ \\ (-1)R_4 \end{array} \\
 & \sim \left[ \begin{array}{cccc|cccc} 1 & -1 & 3 & -1 & 1 & 0 & 0 & 0 \\ 0 & 1 & -2 & 3 & -1 & 1/2 & 0 & 0 \\ 0 & 0 & 3 & -2 & -1 & -1 & 1 & 0 \\ 0 & 0 & 0 & 1 & -2 & 3/2 & 0 & -1 \end{array} \right] \begin{array}{l} R_1 + R_4 \\ R_2 - 3R_4 \\ R_3 + 2R_4 \end{array} \sim \left[ \begin{array}{cccc|cccc} 1 & -1 & 3 & 0 & -1 & 3/2 & 0 & -1 \\ 0 & 1 & -2 & 0 & 5 & -4 & 0 & 3 \\ 0 & 0 & 1 & 0 & -5 & 2/3 & 1/3 & -2/3 \\ 0 & 0 & 0 & 1 & -2 & 3/2 & 0 & -1 \end{array} \right] \begin{array}{l} \\ \\ \frac{1}{3}R_3 \\ \\ \end{array} \\
 & \sim \left[ \begin{array}{cccc|cccc} 1 & -1 & 0 & 0 & -1 & 3/2 & 0 & -1 \\ 0 & 1 & 0 & 0 & 5 & -4 & 0 & 3 \\ 0 & 0 & 1 & 0 & -5 & 2/3 & 1/3 & -2/3 \\ 0 & 0 & 0 & 1 & -2 & 3/2 & 0 & -1 \end{array} \right] \begin{array}{l} R_1 - 3R_3 \\ R_2 + 2R_3 \\ \\ \end{array} \sim \left[ \begin{array}{cccc|cccc} 1 & -1 & 0 & 0 & 4 & -1/2 & -1 & 1 \\ 0 & 1 & 0 & 0 & 5/3 & -8/3 & 2/3 & 5/3 \\ 0 & 0 & 1 & 0 & -5/3 & 2/3 & 1/3 & -2/3 \\ 0 & 0 & 0 & 1 & -2 & 3/2 & 0 & -1 \end{array} \right] \\
 & R_1 + R_2 \sim \left[ \begin{array}{cccc|cccc} 1 & 0 & 0 & 0 & 17/3 & -19/6 & -1/3 & 8/3 \\ 0 & 1 & 0 & 0 & 5/3 & -8/3 & 2/3 & 5/3 \\ 0 & 0 & 1 & 0 & -5/3 & 2/3 & 1/3 & -2/3 \\ 0 & 0 & 0 & 1 & -2 & 3/2 & 0 & -1 \end{array} \right]
 \end{aligned}$$

### The Change of Coordinates Matrix

#### Solution

And so we see that

$$P = Q^{-1} = \begin{bmatrix} 17/3 & -19/6 & -1/3 & 8/3 \\ 5/3 & -8/3 & 2/3 & 5/3 \\ -5/3 & 2/3 & 1/3 & -2/3 \\ -2 & 3/2 & 0 & -1 \end{bmatrix}$$